

1/14

mIL10	19	MPGSALLCCLLLLTGMRISRGQYSREDNNCTHFPVGQSHMLLELR TAF	1
hIL10			
BCRFI		MHSSALLCCLVLLTGVRASPGQGTQSENCTHFPGNL PNLRLRDAF	
		MERRLVVTLQCLVLL--YLAPECGGTD---QCDNFPQ-----MLRDLRDAF	30
		SQVKTFFQTKDQLDNILLTDSLMMQDFKGYLGCQALSEMIQFYLVEMPQAEKHGPEIKEHLNSLG	
		SRVKTFFQMKDQLDNILLKESLLEDFKGYLGCQALSEMIQFYLVEMPQAEQDPDIKAHVNSLG	
		SRVKTFFQTKDEVNDLLKESLLEDFKGYLGCQALSEMIQFYLVEMPQAEQDPPEAKDHYNSLG	31
		EKLKTLRMRLRRCHRFPCENKSKAVEQVKSDFNKLQDQGVYKAMNEFDIFINCI EAYMMIKMKS	95
		ENLKTLRLRLRRCHRFPCENKSKAVEQVKNAFNKLQEKGIYKAMSEFDIFINYIEAYMTMKIRN	
		ENLKTLRLRLRRCHRFPCENKSKAVEQIKNAFNKLQEKGIYKAMSEFDIFINYIEAYMTIKAR	96
			160

Comparison of the predicted amino acid sequences of mIL-10, hIL-10, and BCRFI. Amino acid sequence identities are indicated by vertical lines.

Fig. 1



2/14

COOH TERMINAL PEPTIDE SEQUENCES OF IL-10 INCLUDING  
9 AMINO ACIDS OF PORCINE, HUMAN and VIRUS PROTEINS.

PORCINE IL-10		*		*		*														
PIT	-	A	-	Y	-	M	-	T	-	M	-	K	-	M	-	R	-	K	-	N
HUMAN IL-10				*				*				*								
IT 9302	-	A	-	Y	-	M	-	T	-	M	-	K	-	I	-	R	-	N		
BCRFI IL-10				*				*				*								
VIT	-	A	-	Y	-	M	-	T	-	I	-	K	-	A	-	R	-			

Aminoacids marked with \* represent key position for  
receptor binding and are evolutionary well conserved.  
Compare the IL-10 aminoacid sequence from mouse below.


MOUSE IL-10								*												
MIT	-	A	-	Y	-	M	-	M	-	I	-	K	-	M	-	K	-	S		

This sequence is lacking the homology to the important  
amino acids.

Fig. 2



3/14



IT9302 inhibits spontaneous IL-8 production by purified cultured monocytes. (▲) Indicates the level of IL-8 when using rh IL-10 (100 ng/ml).

Fig. 3



4/14




Fig. 4



5/14

IT9302 stimulated human monocytes




Fig. 5



6/14





Fig. 6



BEST AVAILABLE COPY

7/14

Chemotactic activity of IT9302  
on CD8+ T cells



IT9302 (ng/ml)

Fig. 7



BEST AVAILABLE COPY

8/14

Desensitization of CD8+ T cells by  
IT9302, resulting in unresponsiveness  
of CD8+ T cells towards IL-10 (10 ng/ml) induced chemotaxis




Fig. 8



BEST AVAILABLE COPY

9/14

## Suppression of IL-8 activity by IT9302

Effect of IT9302 on IL-8 (10 ng/ml) mediated chemotaxis.




Fig. 9



BEST AVAILABLE COPY

10/14

IT9302 inhibits MCAF/MCP-1  
induced monocyte chemotaxis





Fig. 10



BEST AVAILABLE COPY

11/14

ECL-Western Blotting of CD4+ T cell cytosolic proteins using a goat anti-human IL-4 antibody.



Stimulation for 3 days with:

1. Control
2. mon.anti-IL-8 antibody WS.4 10  $\mu$ g/ml,
3. rIL-8 100 ng/ml,
4. rIL-10 100 ng/ml,
5. IT9302 10 ng/ml,
6. rIFN gamma 10 ng/ml.


Fig. 11



BEST AVAILABLE COPY

12/14

ECL-Western Blotting of Human Mixed lymphocyte culture cytosolic proteins using a rabbit anti-human TNF- $\alpha$  antibody.



Stimulation for 24 hours with:

1. Control
2. IT9302 10 ng/ml

Fig. 12



13/14




Fig. 13

14/14

Leukocyte count in peripheral blood (rabbits)  
after induction of acute pancreatitis (5% bile acid)




Fig. 14